

Table Caption. Percent species composition of hunter harvested waterfowl in Suisun Marsh (left panel) and California (right panel). Hunter harvest data from U.S. Fish and Wildlife Service's parts survey. Table from Ackerman et al. 2014.

Ackerman, JT, MP Herzog, GS Yarris, ML Casazza, E Burns, and JM Eadie. 2014. Chapter 5: Waterfowl ecology and management. Pages 103-132 and maps 10 and 11 in Moyle, PB, A Manfree, and PL Fiedler (editors): Suisun Marsh: Ecological History and Possible Futures. University of California Press: Berkeley, California. 239 pages.

Species ¹	Hunter Harvest													
	Suisun Marsh							California						
	1953-1960	1961-1969	1970-1979	1980-1989	1990-1999	2000-2009	2010-2011	1953-1960	1961-1969	1970-1979	1980-1989	1990-1999	2000-2009	2010-2011
Northern Pintail	na	46.6%	50.7%	28.3%	13.1%	8.6%	9.3%	na	31.6%	35.9%	22.5%	11.9%	9.1%	14.2%
American Wigeon	na	13.4%	12.2%	13.0%	16.0%	14.4%	13.5%	na	12.8%	11.1%	10.5%	12.2%	13.7%	13.2%
Northern Shoveler	na	11.4%	10.6%	11.4%	14.6%	12.9%	19.9%	na	9.1%	8.4%	9.5%	11.0%	12.2%	12.9%
Mallard	na	10.6%	8.3%	22.0%	23.7%	25.4%	27.7%	na	18.0%	15.5%	23.1%	26.1%	23.5%	19.4%
American Green-winged Teal	na	9.6%	11.1%	14.1%	19.0%	19.7%	17.6%	na	15.9%	17.7%	19.3%	23.9%	24.4%	23.0%
Gadwall	na	1.4%	1.1%	3.5%	5.2%	6.6%	3.7%	na	2.7%	2.4%	4.7%	7.0%	7.8%	7.3%
Greater White-Fronted Goose	na	1.1%	0.6%	0.4%	0.2%	0.9%	1.4%	na	3.2%	2.2%	1.2%	1.5%	3.6%	4.0%
Canada Goose	na	1.1%	0.8%	0.3%	0.5%	2.1%	0.7%	na	4.2%	3.8%	3.5%	2.4%	3.3%	4.0%
Ruddy Duck	na	1.0%	0.8%	0.6%	0.1%	0.4%	1.0%	na	1.2%	1.1%	0.7%	0.4%	0.2%	0.7%
Cinnamon Teal	na	0.9%	0.8%	1.4%	1.5%	2.0%	1.0%	na	3.2%	2.9%	3.3%	3.0%	3.4%	2.8%
Scaup	na	0.8%	0.5%	1.2%	2.1%	2.1%	0.6%	na	1.9%	1.4%	1.6%	1.2%	1.4%	0.8%
Canvasback	na	0.7%	1.2%	1.8%	1.1%	0.7%	0.0%	na	0.7%	1.2%	1.1%	1.1%	0.8%	1.0%
Ring-Necked Duck	na	0.4%	0.3%	0.5%	0.4%	0.3%	0.6%	na	0.7%	0.7%	1.4%	1.8%	1.7%	2.2%
Bufflehead	na	0.3%	0.1%	0.4%	0.8%	0.9%	0.8%	na	0.9%	0.4%	0.7%	0.8%	0.8%	0.5%
Common Goldeneye	na	0.2%	0.1%	0.4%	0.8%	1.4%	1.0%	na	0.4%	0.2%	0.3%	0.3%	0.4%	0.3%
Snow & Ross' Goose	na	0.2%	0.2%	0.3%	0.0%	0.3%	0.0%	na	3.9%	4.3%	4.0%	4.1%	4.0%	4.1%
Redhead	na	0.2%	0.1%	0.2%	0.4%	0.2%	0.0%	na	0.4%	0.5%	0.7%	0.7%	0.6%	0.4%
Wood Duck	na	0.1%	0.1%	0.2%	0.4%	0.4%	0.8%	na	1.1%	1.2%	1.9%	2.3%	2.4%	2.0%
Black Brant	na	0.0%	0.0%	0.0%	0.0%	0.3%	0.0%	na	0.5%	0.2%	0.0%	0.3%	0.1%	0.0%
Common Merganser	na	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	na	0.1%	0.1%	0.1%	0.1%	0.1%	0.1%
Surf Scoter	na	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	na	0.1%	0.1%	0.1%	0.0%	0.0%	0.3%

¹ Some species are not differentiated during mid-winter waterfowl surveys, and we have listed the dominant species only. However, canada goose includes lesser, Taverner, dusky, and western canada goose; cackling goose includes aleutian and small cackling goose; American wigeon may include some Eurasian wigeon; scaup includes both lesser scaup and greater scaup; surf scoter may include some black scoter and white-winged scoter; common goldeneye may include some Barrow's goldeneye; common merganser may include some hooded merganser and red-breasted merganser; and eider includes *Somateria spp.*